



ACTIVE
AgriScience
activeagriscience.com



3 - 19 - 7

In crop
growth
promoter

N-P-K + micronutrients liquid fertiliser solution in an easy to absorb form.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. **DIRECTIONS FOR USE:** Apply at herbicide timing as a foliar spray using 2.5 L / ha with a minimum of 100 L of water / ha.. Allow at least 3-4 weeks between applications. **Wheat and oats:** apply at BBCH 13-15 (3-5 leaf stage). **Canola:** apply at BBCH 12-14 (2-4 leaf stage). **Soybean:** apply at V1-V2 (first-second trifoliolate). **Peas, lentils, and other pulse crops, potato, corn:** apply at the 4-5 leaf stage. **Flax:** apply when plants have 3 sets of true leaves. Spray early morning or late afternoon when the sun is lower in the sky. Do not apply when air temperatures are above 29°C. Avoid spraying on windy days. Use this product on the basis of soil and tissue analysis in accordance with recommendations of a qualified person or institution or apply according to recommendations in your approved nutrient program. This product may be tank mixed with pesticides. In some cases, tank mixing with pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact the company responsible for the pest control product before applying any tank mix. **COMPATIBILITY:** This product is compatible with most pesticides and fertilisers. If compatibility is uncertain, conduct a jar test prior to use. **Add tank mix partners in the following order:** Water, Agrochemical, Active BUILD™. Keep agitator running while mixing and spraying. **WARRANTY AND CONDITIONS OF SALE:** Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied.



DANGER HAZARD STATEMENTS: MAY CAUSE EYE AND SKIN IRRITATION. CONTAINS BORON: MAY DAMAGE FERTILITY OR THE UNBORN CHILD.

CAUTION: THIS FERTILISER CONTAINS BORON, MANGANESE, MOLYBDENUM, AND ZINC AND SHOULD ONLY BE USED AS RECOMMENDED. IT MAY PROVE HARMFUL WHEN MISUSED. Excessive application of molybdenum to pasture can be harmful to stock animals. Plant levels of molybdenum can be high for up to 4 weeks after application. It is advisable to keep stock animals off areas to which this product has been applied for 4 weeks. **PRECAUTIONARY STATEMENTS:** Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear chemical resistant gloves, coveralls, goggles and face protection. If mists are present and ventilation is absent or poor, use an air purifying respirator. Do not eat, drink or smoke when using this product. Wash up thoroughly before eating, drinking, smoking and leaving work. **FIRST AID: IF SWALLOWED:** Rinse mouth. Do NOT induce vomiting. Seek medical attention if you feel unwell. **IF ON SKIN:** Immediately take off all contaminated clothing. Rinse skin with water (or shower). Wash contaminated clothing before reuse. Seek medical attention if irritation persists. **IF INHALED:** Remove person to fresh air and keep comfortable for breathing. Seek medical attention if you feel unwell. **IF IN EYES:** Immediately flush eyes with a gentle stream of water for 15 minutes while holding the upper and lower eyelids open. Remove contact lenses, if present and easy to do. Seek medical attention if irritation persists. **STORAGE:** KEEP OUT OF REACH OF CHILDREN. Store locked up at a minimum temperature of 8°C in a dry, well ventilated area away from food or feed storage. DO NOT FREEZE. **DISPOSAL:** Dispose of this product and its container in accordance with Federal, Provincial, and Local regulations.

FOR CHEMICAL EMERGENCY CALL: +61 1800 951 288
and refer to SDS (scan QR code or visit
activeagri.com/active-build).



Manufactured by Active AgriScience Australia
807/893 Canning Highway
Mt Pleasant WA 6153

ANALYSIS (W/V%):			
Total Nitrogen (N).....	2.9%	Boron (B)(actual)	0.4%
Phosphorous (P)	19.3%	Manganese (Mn)(actual).....	1.5%
Potassium (K)	7.3%	Molybdenum (Mo)(actual)	0.2%
Sulfur (S)	0.9%	Zinc (Zn)(actual)	3.4%

Active AgriScience DISCLAIMER: Please read this limitation carefully as it impacts your legal rights. It is understood and agreed that Active AgriScience Pty Ltd ("Active") does not guarantee that use of this Product will yield any specific results. Active's legal liability, and that of its employees or agents, arising from use of this product shall be limited to the cost paid for the product regardless of whether any loss arose from Active's own negligence or breach of contract or any other cause. Under no circumstances shall Active be liable, beyond the cost paid for the product, for direct, consequential, incidental, or special damages, including, but not limited to, damage or destruction of a crop, reduced yield from any crop, or contamination of any property. Use or sale of this product confirms acceptance of this limitation of legal liability.

activeagri.com/active-build



SCAN ME
for more info